Emerald Ash Borer

Homeowner Guide to Insecticide Selection, Use, and Environmental Protection



What is the Risk to Your Tree(s)?

Trees need to be protected with insecticides when emerald ash borer (EAB) populations are abundant enough in an area to cause significant injury in the next year or two. Deciding when to use insecticides to protect trees from EAB depends on the EAB abundance in the area and how much risk you are willing to tolerate regarding the tree or trees in question. To learn about EAB abundance in your area:

 Check the Minnesota Department of Agriculture's (MDA) Interactive EAB Survey Map for the locations of known infestations in Minnesota at: www.mda.state.mn.us/eab



- Check with your local government or the MDA about monitoring efforts in your area.
- Learn how to recognize symptoms of EAB infestation and keep an eye on the trees in your neighborhood.

Treatment Will Likely Need to be Repeated

Insecticides only work while they are present in the tree, and products are generally labeled for one to two years of effectiveness. Treatment will need to be repeated to protect trees if EAB are still abundant enough in an area to cause significant injury when the effective period ends. EAB could remain at damaging levels in an area for several treatment cycles or return to damaging levels in the future.

Remove and Replace Ash Trees on Your Property

It may be more cost effective to replace a small or struggling ash tree than to pay the cost of ongoing treatments. In addition, trees in poor health are not likely to respond well to treatments. Do not treat trees showing more than 50 percent canopy decline; these ash trees are unlikely to recover even if treated. Check with a certified arborist or licensed professional pesticide applicator to compare costs and benefits of treatment versus removal.

Check Your Calendar – Timing is Everything

To ensure the insecticide is distributed throughout the tree to control EAB, soil-applied insecticides available to homeowners are most effective when applied mid-to late spring or in mid-fall. Do not apply products when the soil is frozen. Refer to the table on page 2 for more information on the timing of different insecticide options. Weather conditions or other factors may alter these recommendations. Read pesticide labels carefully. **The label is the law.**

Contact a Certified Arborist or City Forester to Evaluate Treatment Options

When contacting a professional, consider the following:

- Your city or township may have requirements or restrictions related to the treatment of EAB, especially for ash trees located on
 city property, which may include trees planted on or near boulevards or sidewalks. Check with your city's park and recreation or
 forestry departments.
- If hiring a professional, ask to see their licensure as a Commercial Pesticide Applicator for category "E: Turf and Ornamentals."

Treatment of Large Ash Trees

Treatment options available to homeowners for treating large ash trees require carefully following application instructions to avoid unintended environmental impacts. In general, consider having large ash trees (those greater than 48 inches in circumference, as measured 4½ feet above ground level) treated by a professional. Check specific pesticide product labels for tree size requirements.







Follow These Recommendations to Protect Water Quality

The MDA conducted a special registration review of EAB insecticides in 2011. The review concluded that insecticides commonly used to control EAB are not likely to result in unreasonable risks to human health or the environment when used according to label directions.


In some cases, the following recommendations may refer to mandatory label use requirements:

Generally, professionally applied trunk injections have fewer concerns for water quality.

For soil-applied products and bark sprays, the following considerations are important:

- Properly prepare application site and apply product according to label instructions. Consider avoiding use
 within 25 feet of water bodies. This includes streams, lakes, ponds, wetlands, or conduits to surface water or
 groundwater such as street curbs, storm drains, sumps, or well heads.
- Do not apply when heavy rainfall is expected within 24 hours of the planned treatment.

Situations that could cause soil-applied insecticides to wash away or leach, potentially contaminating water resources



Making Sense of EAB Insecticide Labels

The label is the law! Always read, understand, and follow the label directions for the specific insecticide product being used.

Selecting and Using an Insecticide:


- Look for products marketed to control emerald ash borer (see Products Table on page 2).
- Read the Environmental Hazard Statements on the insecticide label.
- To facilitate uptake, soil-applied insecticides should be applied when the soil is moist but not saturated or excessively dry.
- For soil drenches, remove or rake any mulch or dead leaves, then pour the insecticide solution directly onto soil.
- Do not allow children and pets to re-enter treatment area until sprays or drenches have dried.
- Store insecticides where children cannot reach them.
- Many homeowner products only allow one soil drench application per year.
- Some products have maximum per acre use limits. Do not apply more than the approved labeled rate.

Disposing an Insecticide:

- When using the last of a liquid insecticide, triple-rinse the container before disposal, then apply the rinse water as you would apply the insecticide.
- Unusable and unwanted insecticides must be disposed of according to the label directions, or at a county household hazardous waste disposal event.
- It is illegal to bury or burn an insecticide.

Measure Your Ash Trees

It is important to measure your ash tree to determine the appropriate treatment and to accurately follow the use instructions on insecticide labels.



trunk	diameter	=	trunk	circumference	X 0.32

CONVERSION TABLE Tree Measurements at 4½ Feet Above Ground Level					
Circumference – Inches	Diameter at Breast Height (DBH) – Inches				
15	5				
20	6				
25	8				
30	9.5				
35	11				
40	13				
45	14				
> 48 inches	> 15 inches				
In general large trees are best treeted by a					

In general, large trees are best treated by a professional. For exceptions, check specific pesticide product labels.

Additional Resources

Emerald Ash Borer Resources

- Minnesota Department of Agriculture www.mda.state.mn.us/eab
- EAB Multi-State Site www.emeraldashborer.info
- University of Minnesota Extension www.extension.umn.edu/garden/insects/find/emerald-ash-borer/
- USDA-APHIS www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b
- Minnesota Department of Natural Resources www.mndnr.gov/invasives/terrestrialanimals/eab

Pesticide Disposal

For more information about disposing of unusable or unwanted pesticides, visit MDA at www.mda.state.mn.us/chemicals/spills/wastepesticides/schedule.aspx or 651-201-6562.

Reporting and Investigating Pesticide Spills and Misuse

Complaints can be reported to the Minnesota Duty Officer at 800-422-0798.

The MDA staff will evaluate the information provided and if a pesticide misuse is suspected, an the MDA inspector will be assigned. More information is available at www.mda.state.mn.us/chemicals/pesticides/complaints.aspx